

ATPase activity of the microvillar 110 kDa polypeptide-calmodulin complex is activated in Mg^{2+} and inhibited in K^{+} -EDTA by F-actin

Janet Krizek, Lynne M. Coluccio and Anthony Bretscher

Section of Biochemistry, Molecular and Cell Biology, Wing Hall, Cornell University, Ithaca, NY 14853, USA

Received 28 October 1987

Highly purified microvillar 110 kDa polypeptide-calmodulin (110K-cam) complex was confirmed to have ATPase activities characteristic of a myosin. The effect of F-actin on these activities was investigated. The Mg^{2+} -ATPase is activated about 2-fold by F-actin in a dose-dependent fashion, whereas the K^{+} -EDTA-ATPase is inhibited by >90% by F-actin. These data provide evidence for a functional relationship between the ATPase activity of 110K-cam and its interaction with F-actin. They also extend the similarities between 110K-cam and myosin. The results suggest that higher cells contain in addition to myosin a second class of myosin-like molecules represented by 110K-cam.

Calmodulin; Myosin; Actin; Microvillus; Cytoskeleton; ATPase

1. INTRODUCTION

Regularly spaced lateral bridges attach the core bundle of actin filaments of intestinal microvilli to the plasma membrane [1,2]. Recently, evidence has been obtained to suggest that these bridges are comprised of a 110 kDa polypeptide-calmodulin complex (110K-cam) [3–8]. The purified complex shares some properties with muscle and non-muscle myosins, although these myosins typically consist of two heavy chains of ~200 kDa and four light chains of ~18 kDa. Similarities between 110K-cam and the myosins include: (i) high K^{+} -EDTA and Ca^{2+} -ATPase activities and low Mg^{2+} -ATPase activity [6,9]; (ii) saturable binding to F-actin in an ATP-reversible manner [7]; and (iii) the ability to ‘decorate’ F-actin to give rise to an arrowhead appearance [7]. Moreover, as with

myosins, both the ATPase and F-actin binding activities reside on the heavy polypeptide chain and are independent of the calmodulin ‘light chains’ [8]. The Mg^{2+} -ATPase activity has been reported not to be activated by F-actin [6] or to be modestly activated [9]. 110K-cam has not yet been demonstrated to couple ATP hydrolysis with mechanical work.

We recently described an improved method for the purification of native 110K-cam and explored its interaction with F-actin [7]. Here, we investigate the ATPase activity of 110K-cam in relation to its interaction with F-actin. We show that its ATPase activity, like that of myosin, is enhanced by F-actin in the presence of Mg^{2+} and inhibited by F-actin in its absence.

2. MATERIALS AND METHODS

Microvillar 110K-cam was purified as described [7]. Brush border myosin was purified according to

Correspondence address: J. Krizek, Section of Biochemistry, Molecular and Cell Biology, Wing Hall, Cornell University, Ithaca, NY 14853, USA

Table 1

Specific ATPase activities of 110K-cam and brush border myosin

	1 mM EDTA, 1 mM Ca^{2+} 5 mM Mg^{2+} 1 mM EGTA ^a		
110K-cam			
50 mM KCl	502	357	14
0.5 M KCl	211	105	19
Myosin			
0.5 M KCl	625	382	10

^a All assays were performed in ATPase buffer with additions to give the final concentrations indicated. Values are expressed in nmol P_i /mg protein per min

a new procedure, adapted from [6]. Rabbit skeletal muscle actin was purified [10] and gel filtered [11] before use. ATPase assays were performed [12] in 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, pH 8.0, with appropriate additions as indicated, at 37°C. The assay was found to be linear with time up to 30 min (the longest time point used). All the assays were repeated on a number of different 110K-cam preparations and representative values are presented in section 3. Gel electrophoresis in the presence of SDS was as described [7,13]. Protein was determined colorimetrically [14] using bovine serum albumin as a standard.




Fig.1. 7.5%/15% SDS-polyacrylamide gel of purified microvillar 110K-cam (A), and purified brush border myosin (B). Molecular mass standards (in kDa) are indicated in lane C.




Fig.2. Effect of KCl concentration on the ATPase activities of 110K-cam. The curves represent the activities in the presence of 5 mM MgCl_2 , 5 mM CaCl_2 and 1 mM EDTA, 1 mM EGTA.

3. RESULTS

The typical ATPase activities of purified 110K-cam are listed in table 1, together with results obtained using brush border myosin. The purity of a 110K-cam preparation and brush border myosin as assessed by SDS gel electrophoresis is shown in fig.1. Since the activities of myosins are typically measured in 0.6 M KCl to inhibit filament forma-




Fig.3. Effect of F-actin on the Mg^{2+} -ATPase activity of 110K-cam. The assays contained 100 $\mu\text{g}/\text{ml}$ 110K-cam and the indicated amounts of actin. The very low ATPase activity of F-actin alone was subtracted from the values obtained.




Fig.4. Effect of F-actin on the ATPase activity of 110K-cam in the presence of 1 mM EDTA, 1 mM EGTA. The assays contained 20 μ g/ml 110K-cam and the indicated amounts of actin. The very low ATPase activity of F-actin alone was subtracted from the values obtained.

tion, we explored the effect of KCl concentration on the ATPase activities of 110K-cam and found that the K^+ -EDTA/EGTA and Ca^{2+} -ATPases were inhibited by increasing KCl concentrations, whereas Mg^{2+} -ATPase was slightly enhanced (fig.2). Since 110K-cam does not aggregate in low salt [7] and these conditions are more representative of a normal physiological state, all further ATPase measurements on 110K-cam were made at relatively low KCl concentrations.

Fig.3 shows the effects of F-actin on Mg^{2+} -ATPase of 110K-cam. F-Actin activation was invariably found, although the degree of activation varied from about 2- to 3-fold from one 110K-cam preparation to another. The Ca^{2+} -ATPase activity was either unaffected by the presence of F-actin, or slightly reduced (not shown).

By contrast, the K^+ -EDTA/EGTA activity was invariably found to be inhibited by ~90% in the presence of F-actin (fig.4). In a similar manner, the K^+ -EDTA/EGTA ATPase of 0.1 mg/ml myosin was reduced from 625 to 14 nmol P_i /mg per min by the presence of 0.2 mg/ml F-actin.

4. DISCUSSION

In this report we confirm that microvillar 110K-

cam is an ATPase with the enzymatic characteristics of a myosin. The specific activities are somewhat higher than those reported by others [6,9], but show the same ratio between the K^+ -EDTA/EGTA, Ca^{2+} - and Mg^{2+} -ATPase activities. The higher activities may be due to the greater purity of the preparation and the lack of proteolytic products. We also show that 110K-cam exhibits two enzymatic characteristics of a myosin: an Mg^{2+} -ATPase that can be modestly activated by F-actin, and a K^+ -EDTA/EGTA ATPase activity that is inhibited by F-actin. The reason for the variable activation of the Mg^{2+} -ATPase by F-actin is not yet clear. One possibility is that, like myosin, it depends on the degree of phosphorylation [15]. In this regard, it is of interest to note that the 110 kDa polypeptide is a substrate for a kinase present in microvillar preparations [5].

The present data together with earlier results [5], may give some insight into the interaction between 110K-cam and F-actin. Dissociation of 110K-cam from F-actin requires both ATP (or other nucleoside triphosphates) and a divalent cation [5]. Thus, in the presence of K^+ -EDTA/EGTA and ATP, 110K-cam remains bound to F-actin, but dissociates in Mg^{2+} and ATP. The data are consistent with a model in which 110K-cam associates repeatedly with F-actin in Mg^{2+} and ATP, and thereby increases 110K-cam's Mg^{2+} -ATPase activity, whereas in K^+ -EDTA/EGTA 110K-cam is bound to F-actin which inhibits this ATPase activity. These results provide further evidence for a functional correlation between the ATPase activity of 110K-cam and its ability to bind F-actin in an ATP-reversible manner. Further, they suggest that hydrolysis of Mg^{2+} -ATP may be required in the binding/dissociation cycle of 110K-cam and F-actin, in a manner similar to the cross-bridge cycle between myosin heads and F-actin.

The results add substantially to the emerging biochemical similarities between 110K-cam and bona fide myosins and suggest that there may be at least two distinct classes of myosin-like molecules in higher non-muscle cells.

ACKNOWLEDGEMENTS

This work was supported by National Institutes of Health grants GM36652 to A.B. and AI 24560 to L.M.C.

REFERENCES

- [1] Brunser, O. and Luft, J.H. (1970) *J. Ultrastruct. Res.* 31, 291–311.
- [2] Mukherjee, T.M. and Staehelin, L.A. (1971) *J. Cell Sci.* 8, 573–599.
- [3] Matsudaira, P.T. and Burgess, D.R. (1979) *J. Cell Biol.* 83, 667–673.
- [4] Howe, C.L. and Mooseker, M.S. (1983) *J. Cell Biol.* 97, 974–985.
- [5] Verner, K. and Bretscher, A. (1985) *J. Cell Biol.* 100, 1455–1465.
- [6] Collins, J.H. and Borysenko, C.W. (1984) *J. Biol. Chem.* 259, 14128–14135.
- [7] Coluccio, L.M. and Bretscher, A. (1987) *J. Cell Biol.* 105, 325–333.
- [8] Coluccio, L.M. and Bretscher, A. (1987) submitted.
- [9] Conzelman, K.A. and Mooseker, M.S. (1987) *J. Cell Biol.* 105, 313–324.
- [10] Spudich, J.A. and Watt, S. (1971) *J. Biol. Chem.* 246, 4866–4871.
- [11] MacLean-Fletcher, S. and Pollard, T.D. (1980) *Biochem. Biophys. Res. Commun.* 96, 18–27.
- [12] Pollard, T.D. (1982) *Methods Cell Biol.* 24, 333–370.
- [13] Laemmli, V. (1971) *Nature* 227, 680–685.
- [14] Bradford, M.M. (1976) *Anal. Biochem.* 72, 248–254.
- [15] Sobieszek, A. and Small, J.V. (1977) *J. Mol. Biol.* 112, 559–576.